

KEYWORD SEARCH

GO

OUR PRODUCTS

STORE LOCATOR

ABOUT US

FAQs

LINKS

CONTACT US


B.F. ASCHER & CO. INC.
pharmaceuticals
*Makers of Consumer Health Care
Products for Better Living.*


Mobigesic Tablets

- Provides pain relief and mild relaxation
- Relieves pain associated with headache, toothache, menstrual cramps
- Less stomach irritation than aspirin products

[LOCATE THIS PRODUCT IN A STORE NEAR YOU](#)
[Click to return to the **PRODUCT CATEGORY PAGE**](#)

Provides temporary relief of minor aches and pains associated with back and muscle ache

Help your aching back and muscles with Mobigesic tablets. Mobigesic provides temporary relief of minor aches and pains associated with back ache and muscle ache. Containing both a pain reliever (magnesium salicylate 325 mg) and an antihistamine (phenyltoloxamine citrate 30 mg), Mobigesic provides pain relief and mild relaxation to help you get past your pain and return to your active life. Mobigesic causes less irritation to your stomach than aspirin products. Also effective in relieving pain associated with headache, toothache, and menstrual cramps.

INGREDIENTS

Active Ingredients Per Tablet: Magnesium salicylate tetrahydrate 404 mg (Equivalent to 325 mg of magnesium salicylate anhydrous.)
Phenyltoloxamine citrate 30 mg

Inactive Ingredients: magnesium stearate, microcrystalline cellulose, povidone and silicon dioxide.

DIRECTIONS

Adults: Drink a full glass of water with each dose. Take 1 or 2 tablets every 4 hours while symptoms persist. Do not take more than 10 tablets in 24 hours.

USES

For the temporary relief of minor aches and pains associated with:

- Backache and muscular aches
- Back pain due to muscle strain or spasm
- Muscle stiffness

Purposes: Analgesic and antihistamine (analgesic adjuvant)

WARNINGS

Do not take this product if you are taking a prescription drug for anticoagulation (thinning of the blood), diabetes, gout or arthritis unless directed by a physician.

Do not take for more than 10 days unless directed by a physician.

May cause drowsiness. Alcohol, sedatives, and tranquilizers may increase the drowsiness effect. Use caution when driving a motor vehicle or operating machinery.

Children and teenagers who have or are recovering from chicken pox, flu symptoms, or flu should NOT use this product before a physician is consulted about Reye's syndrome, a rare but serious illness.

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your physician whether you should take magnesium salicylate or other pain relievers/fever reducers. Magnesium salicylate may cause stomach bleeding.

AVAILABILITY

In bottles of 18, 50 and 100 Mobigesic Tablets.

[HOME](#)

[DISCLAIMER / PRIVACY POLICY](#)